Meadow Feeds Broiler Product Range





meadowfeeds.co.za

INDEX

Product		Page
Broiler Feed	ł	
V 16371	Classic Broiler Starter	1
V 17142	Classic Broiler Grower	2
V 16419	Classic Broiler Finisher	3
V 16420	Classic Broiler Post-finisher	4
V 16640	Select Broiler Pre-starter	5
V 10757	Select Broiler Starter	6
V 10754	Select Broiler Grower	7
V 10761	Select Broiler Finisher	8
V 17488	Select Broiler Post-finisher	9
V 17767	Prime Broiler Starter	10
V 17768	Prime Broiler Grower	11
V 17769	Prime Broiler Finisher	12
V 17810	Prime Broiler Post-finisher	13
V 19502	Budget Broiler Starter	14
V 19501	Budget Broiler Grower	15
V 19506	Budget Broiler Finisher	16
V 18343	Broiler Maintenance	17





Form: Pellets / Mash / Crumbles

Description

The Classic Broiler range has been specifically designed for medium-sized producers who produce broilers between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Classic feeding programme was formulated to optimise performance for medium-scale enterprises, considering environmental conditions and average-to-low stocking density.

Feeding Recommendations

Feed broilers from day old to 18 days of age. The approximate feed consumption is 800 g per bird. Crumbles are fed to improve early feed intake and bird development.

Expected Intakes

The approximate feed consumption is 800 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	200	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	8	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	6	g/kg	Min
Total lysine	12	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

	71		,	
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Pellets / Mash / Crumbles

Description

The Classic Broiler range has been specifically designed for medium-sized producers who produce broilers between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Classic feeding programme was formulated to optimise performance for medium-scale enterprises, considering environmental conditions and average-to-low stocking density.

Feeding Recommendations

The grower ration should be fed to broilers from 19–28 days of age. The approximate feed consumption should be 1 200 g per bird. The ration may be fed through to term, but this is not advisable.

Expected Intakes

The approximate feed consumption should be 1 200 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	180	g/kg	Min
Fat	25	g/kg	Min
Fibre	60	g/kg	Max
Moisture	120	g/kg	Max
Calcium	7	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5.5	g/kg	Min
Total lysine	10	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

	· 1		,	
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



+27 21 807 8700
+27 28 551 1950
+27 41 402 5000

- +260 967 790 404



Class: Broiler Finisher Feed

Form: Pellets

Description

The Classic Broiler range has been specifically designed for medium-sized producers who produce broilers between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Classic feeding programme was formulated to optimise performance for medium-scale enterprises, considering environmental conditions and average-to-low stocking density.

Feeding Recommendations

Broilers should be fed the finisher diet from 29–32 days. The approximate feed consumption should be 900 g per bird. If birds are marketed at an older age (beyond 35 days), the amount of finisher fed should be adjusted to allow for a 5-day withdrawal feed to be used, such as a post-finisher.

Expected Intakes

The approximate feed consumption should be 900 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	70	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Registration No: V 16420 Act 36 of 1947

Class: Broiler Post-Finisher Feed

Form: Pellets / Mash / Crumbles

Description

The Classic Broiler range has been specifically designed for medium-sized producers who produce broilers between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Classic feeding programme was formulated to optimise performance for medium-scale enterprises, considering environmental conditions and average-to-low stocking density.

Feeding Recommendations

Feed 5 days prior to sale or slaughter for human consumption.

Expected Intakes

The approximate feed consumption should be 500 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	60	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Mash / Crumbles

Description

The Select Broiler range has been specially designed for farmers who produce broilers to slaughter at 5 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Select programme was formulated to optimise performance for medium- to large-scale enterprises, considering environmental conditions and increased stocking density.

Feeding Recommendations

Feed broilers from day old to 9 days of age. The approximate feed consumption is 200 g per bird. Crumbles are fed to improve early feed intake and bird development.

Expected Intakes

The approximate feed consumption is 200 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	210	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	8	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	6	g/kg	Min
Total lysine	13	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Mash / Crumbles

Description

The Select Broiler range has been specially designed for farmers who produce broilers to slaughter at 5 weeks of age. This diet specification ensures an optimum dressed bird. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Select programme was formulated to optimise performance for medium- to large-scale enterprises, considering environmental conditions and increased stocking density.

Feeding Recommendations

Feed broilers from 10–18 days of age. The approximate feed consumption is 600 g per bird. Crumbles are fed to improve early feed intake and bird development.

Expected Intakes

The approximate feed consumption is 600 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	200	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	8	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	6	g/kg	Min
Total lysine	12	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Pellets / Mash / Crumbles

Description

The Select Broiler range has been specially designed for farmers who produce broilers to slaughter at 5 weeks of age. This diet specification ensures an optimum dressed bird. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Select programme was formulated to optimise performance for medium- to large-scale enterprises, considering environmental conditions and increased stocking density.

Feeding Recommendations

The grower ration should be fed to broilers from 19-28 days of age. The approximate feed consumption should be 1 200 g per bird.

Expected Intakes

The approximate feed consumption is 1 200 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Class: Broiler Finisher Feed

Form: Pellets

Description

The Select Broiler range has been specially designed for farmers who produce broilers to slaughter at 5 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Select programme was formulated to optimise performance for medium- to large-scale enterprises, considering environmental conditions and increased stocking density.

Feeding Recommendations

The finisher ration should be fed to broilers from 29-32 days of age. The approximate feed consumption should be 850 g per bird. If birds are marketed at an older age (beyond 35 days), the amount of finisher fed should be adjusted to allow for a 5-day withdrawal feed to be used, such as a post-finisher.

Expected Intakes

The approximate feed consumption should be 850 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

	<i>'</i> 1		,	
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia

- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000 - +260 967 790 404





Class: Broiler Post-Finisher Feed

Form: Pellets / Mash / Crumbles

Description

The Select Broiler range has been specially designed for farmers who produce broilers to slaughter at 5 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Select programme was formulated to optimise performance for medium- to large-scale enterprises, considering environmental conditions and increased stocking density.

Feeding Recommendations

Feed 5 days prior to sale or slaughter for human consumption.

Expected Intakes

The approximate feed consumption should be 500 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	60	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

			,	
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia







Form: Mash / Crumbles

Description

The Prime Broiler range has been specially designed for small producers with performance as a benchmark, who market birds between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Prime feeding programme was formulated to optimise performance for a small to medium enterprise, considering environmental conditions and stocking density.

Feeding Recommendations

Feed broilers from day old to 18 days of age. The approximate feed consumption is 800 g per bird. Crumbles are fed to improve early feed intake and bird development.

Expected Intakes

The approximate feed consumption is 800 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	200	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	8	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	6	g/kg	Min
Total lysine	12	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

	<i>,</i> 1		,	
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Mash / Crumbles

Description

The Prime Broiler range has been specially designed for small producers with performance as a benchmark, who market birds between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Prime feeding programme was formulated to optimise performance for a small to medium enterprise, considering environmental conditions and stocking density.

Feeding Recommendations

The grower ration should be fed to broilers from 19–28 days of age. The approximate feed consumption should be 1 200 g per bird. The ration may be fed through to term, but this is not advisable.

Expected Intakes

The approximate feed consumption is 1 200 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	180	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	7	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5.5	g/kg	Min
Total lysine	10	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	· 1	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



+27 21 807 8700
+27 28 551 1950
+27 41 402 5000

- +260 967 790 404



Registration No: V 17769 Act 36 of 1947

Class: Broiler Finisher Feed

Form: Pellets

Description

The Prime Broiler range has been specially designed for small producers with performance as a benchmark, who market birds between 6 and 8 weeks of age. The rations in this range, except the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Prime feeding programme was formulated to optimise performance for a small to medium enterprise, considering environmental conditions and stocking density.

Feeding Recommendations

The finisher ration should be fed to broilers from 29-32 days of age. The approximate feed consumption should be 950 g per bird. If birds are marketed at an older age (beyond 35 days), the amount of finisher fed should be adjusted to allow for a 5-day withdrawal feed to be used, such as a post-finisher.

Expected Intakes

The approximate feed consumption should be 950 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Class: Broiler Post-Finisher Feed

Form: Pellets

Description

The Prime Broiler range has been specially designed for small producers with performance as a benchmark, who market birds between 6 and 8 weeks of age. The rations in this range, with the exception of the post-finisher, include a coccidiostat and a growth enhancer.

Benefits

The Prime feeding programme was formulated to optimise performance for a small to medium enterprise, considering environmental conditions and stocking density.

Feeding Recommendations

Feed 5 days prior to sale or slaughter for human consumption.

Expected Intakes

The approximate feed consumption should be 500 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fat	70	g/kg	Max
Fibre	60	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

	-	07 11 000 7100		- .
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia







Form: Pellets / Mash / Crumbles

Description

The Budget Broiler range has been specifically designed for small producers who market birds between 6 and 8 weeks of age. The rations in this range include a coccidiostat and a growth enhancer.

Benefits

The Budget feeding programme was formulated to optimise performance for small enterprises, considering environmental conditions and stocking density.

Feeding Recommendations

Feed broilers from day old to 18 days of age. The approximate feed consumption is 800-1 000 g per bird. Crumbles are fed to improve early feed intake and bird development.

Expected Intakes

The approximate feed consumption is 800–1 000 g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	200	g/kg	Min
Fat	25	g/kg	Min
Fibre	50	g/kg	Max
Moisture	120	g/kg	Max
Calcium	8	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	6	g/kg	Min
Total lysine	12	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein Standerton	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
KWA-ZULU NATAL	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Form: Pellets / Mash / Crumbles

Description

The Budget Broiler range has been specifically designed for small producers who market birds between 6 and 8 weeks of age. The rations in this range include a coccidiostat and a growth enhancer.

Benefits

The Budget feeding programme was formulated to optimise performance for small enterprises, considering environmental conditions and stocking density.

Feeding Recommendations

The grower ration should be fed to broilers from 19-32 days of age. The approximate feed consumption should be 1 300–1 500 g per bird. The ration may be fed through to term depending on marketing strategies, but this is not advisable.

Expected Intakes

The approximate feed consumption should be 1 300-1 500 g per bird

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	180	g/kg	Min
Fat	25	g/kg	Min
Fibre	60	g/kg	Max
Moisture	120	g/kg	Max
Calcium	7	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5.5	g/kg	Min
Total lysine	10	g/kg	Min

For further technical assistance, please contact a technical advisor in your area.

			,		
CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl	
	Standerton	- +27 17 200 0272		Ladismith	
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha	
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia	



- +27 21 807 8700 - +27 28 551 1950 - +27 41 402 5000

- +260 967 790 404



Class: Broiler Finisher Feed

Form: Pellets / Mash / Crumbles

Description

The Budget Broiler range has been specifically designed for small producers who market birds between 6 and 8 weeks of age. The rations in this range include a coccidiostat and a growth enhancer.

Benefits

The Budget feeding programme was formulated to optimise performance for small enterprises, considering environmental conditions and stocking density.

Feeding Recommendations

The finisher feed should be fed to broilers from 32–42 days. The approximate feed consumption should be 1 500–1 600 g per bird.

Expected Intakes

The approximate feed consumption should be 1 500–1 600g per bird.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	160	g/kg	Min
Fat	25	g/kg	Min
Fibre	70	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	9	g/kg	Min

CENTRAL REGION	Dandfantain	- +27 11 693 5120	WESTERN CAPE	Paarl
CENTRAL REGION			WESTERN CAPE	
	Standerton	- +27 17 200 0272		Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia







Class: Complete Broiler Maintenance Feed

Form: Pellets / Mash / Crumbles

Description

The **Broiler Maintenance** diet has been specially designed for farmers who need to maintain broilers during difficult market conditions for a short period of time.

Benefits

Formulated to maintain body condition for a short period of time at a low cost.

Feeding Recommendations

Feed ad lib to broilers after 5 weeks of age.

Composition

Nutrient	Quantity	Unit	Min/Max
Protein	120	g/kg	Min
Fat	25	g/kg	Min
Fibre	90	g/kg	Max
Moisture	120	g/kg	Max
Calcium	6	g/kg	Min
Calcium	12	g/kg	Max
Phosphorus	5	g/kg	Min
Total lysine	5.5	g/kg	Min

CENTRAL REGION	- 1	- +27 11 693 5120 - +27 17 200 0272	WESTERN CAPE	Paarl Ladismith
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia

