



COMPLETE SHEEP FINISHER FEED 12

Registration No: V 9902 Act 36 of 1947

Class: Complete Sheep Feed
Contains Urea @ 10 g/kg max

Form: Pellets

Description

This ration is a very high-quality complete sheep fattening feed.

Benefits

- Contains the ideal balance between digestible fibre, fat and starch to achieve optimal growth rates.
- Formulated to optimise rumen function.
- Balanced with sufficient by-pass protein.
- Contains the required vitamins, trace- and macro-minerals.

Feeding Recommendations

- Start by feeding 300 g per sheep per day plus roughage.
- Increase gradually until the sheep are on ad-lib feeding after 2 weeks.
- To be fed ad-lib with clean fresh water.
- No additional roughage is needed after successful adaptation.
- Vaccinate for Pulpy Kidney before feeding.
- Offer hay or **Meadow Feeds Drought Cubes 10** ad-lib for 4 days before feeding.

Expected Intakes

Expected feed intake is approximately 3–4% of live body weight.

Composition

Nutrient	Quantity	Unit	Min/Max
*Protein	120	g/kg	Min
Urea	10	g/kg	Max
Fat	25	g/kg	Min
Fat	70	g/kg	Max
Fibre	110	g/kg	Min
Fibre	200	g/kg	Max
Moisture	120	g/kg	Max
Calcium	10	g/kg	Max
Phosphorus	2	g/kg	Min

*23.9 % derived from urea

This product contains NPN sources - see NPN warning on next page

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein - +27 11 693 5120	WESTERN CAPE	Paarl - +27 21 807 8700
	Standerton - +27 17 200 0272		Ladismith - +27 28 551 1950
	Delmas - +27 13 665 5011	EASTERN CAPE	Gqeberha - +27 41 402 5000
KWA-ZULU NATAL	Pietermaritzburg - +27 33 387 2403	AFRICA	Zambia - +260 967 790 404



More than just feed

NPN Warning

This feed may contain feed-grade urean / NPN sources and thus must be mixed and fed strictly according to recommendations:

1. Vinegar is an effective remedy against NPN (non-protein nitrogen) poisoning. Mix with equal amounts of water. Dose half a bottle per calf or large sheep or 2-4 bottles per head of cattle. (1bottle = 750ml)
2. Protect these feeds against rain. NPN is soluble and animals drinking such a solution can be poisoned.
3. Do not feed these feeds indiscriminately with other feed containing NPN. Consult an animal scientist.
4. Mix concentrates thoroughly with the prescribed ingredients.
5. This is a supplement and not a feed. Sufficient grazing and/or roughage must be available at all times.
6. Ensure that feed troughs are always full and give animal's free access to grazing.
7. A salt-phosphate lick must be fed for a minimum of seven days prior to feeding any animal a NPN containing lick.

For further technical assistance, please contact a technical advisor in your area.

CENTRAL REGION	Randfontein	- +27 11 693 5120	WESTERN CAPE	Paarl	- +27 21 807 8700
	Standerton	- +27 17 200 0272		Ladismith	- +27 28 551 1950
	Delmas	- +27 13 665 5011	EASTERN CAPE	Gqeberha	- +27 41 402 5000
KWA-ZULU NATAL	Pietermaritzburg	- +27 33 387 2403	AFRICA	Zambia	- +260 967 790 404



More than just feed